

PLANE DEPTH SEQUENCING

ERIC BAUERFELD

ABSTRACT. Given a plane embedding of a graph with outer cycle C , the *plane depth* of a vertex is its graph distance to C . We use this depth function to organize plane triangulations into a layered combinatorial structure. First, we show that the subgraph induced by each depth level is outerplanar, recovering Baker’s notion of a k -outerplanar graph. We then introduce the *deep embedding* of a maximal planar graph, obtained by inserting a new vertex into each neutral triangular face (including the outer face), and prove that every face of the resulting spherical triangulation is an *up* or *down* triangle. Pairing adjacent triangles across their unique level edge yields a *quadrilateral decomposition* into three combinatorial types: shallow diamonds, deep diamonds, and S quads. Finally, we define a deterministic traversal of this decomposition using four moves — anchor drop, level add, join, and ring completion — under a fixed precedence, and prove that, starting from any boundary deep diamond, the resulting sequence visits every quadrilateral exactly once.

1. MOTIVATION

This paper is one step of a longer programme aimed at 4-coloring maximal planar graphs by inductive local construction. The quadrilateral sequencing is designed so that the deep embedding G' can be built up one quadrilateral at a time, and so that — at least at first glance — each new quadrilateral admits a natural local 4-coloring choice for the vertices it introduces.

Three of the four moves — anchor drop, level add, and join — attach a quadrilateral that introduces one or two new vertices. For each of these moves the new vertex (or vertices) can be assigned a color that respects the local triangle constraints. The fourth move, *ring completion*, introduces no new vertices: it attaches a quadrilateral all four of whose corners are already in the slice. Whether the move succeeds therefore depends on previously chosen colors.

A natural question is whether the local freedom present at the non-ring-completion moves is enough to guarantee that, by the time each ring completion fires, the four corner colors are already compatible with a proper 4-coloring of the new quadrilateral. The remaining sections develop the structural machinery needed to phrase this question precisely; companion commentary (`commentary.tex`) records what is known so far about why this hope is delicate, and an empirical check (`quad_sequence_coloring_check.py`) explores the behavior of a particular online coloring discipline on small triangulations.

2010 *Mathematics Subject Classification*. Primary .

Key words and phrases. plane graph, triangulation, plane depth, deep embedding, quadrilateral decomposition, k -outerplanar graph, canonical sequencing.

2. DEFINITIONS

Definition 2.1. Let G be a graph with a plane embedding, and let C be the outer cycle of that embedding. The *plane depth* of a vertex $v \in V(G)$ relative to the embedding and C is

$$\text{depth}(v) = \min_{u \in V(C)} d(v, u),$$

where $d(v, u)$ denotes the graph distance between v and u in G .

Definition 2.2. An edge $\{u, v\} \in E(G)$ is a *level edge* if $\text{depth}(u) = \text{depth}(v)$.

Definition 2.3. A triangle $\{u, v, w\}$ in G is an *up triangle* if the multiset of depths of its vertices is $\{d, d+1, d+1\}$ for some $d \geq 0$, a *down triangle* if the multiset of depths is $\{d, d, d+1\}$ for some $d \geq 0$, and a *neutral triangle* if the multiset of depths is $\{d, d, d\}$ for some $d \geq 0$.

Remark 2.4. We now relate our terminology to existing terminology, namely k -outerplanar graphs [1]. The following definition and lemma show that the subgraph induced by any single depth level is outerplanar, i.e., 1-outerplanar in the sense of Baker.

Definition 2.5. A plane graph is *outerplanar* if every vertex lies on the outer face. More generally, a plane graph is *k -outerplanar* for $k \geq 1$ if removing all vertices on the outer face yields a $(k-1)$ -outerplanar graph, where every graph on the empty vertex set is 0-outerplanar.

Lemma 2.6. *Let G be a graph with a plane embedding and outer cycle C . For each $d \geq 0$, the subgraph of G induced by $V_d = \{v \in V(G) : \text{depth}(v) = d\}$ is outerplanar.*

Proof. Let $H = G[V_d]$ with the plane embedding inherited from G . It suffices to show every vertex of H lies on the outer face of H .

For $d = 0$, we have $V_0 = C$, so H is outerplanar.

For $d \geq 1$, let U be the open subset of the plane obtained by removing all vertices and edges of H . We show every $v \in V_d$ lies on the boundary of the component U_{out} of U containing the outer face of G .

Since every vertex in $V_{\leq d-1}$ has a shortest path to C passing entirely through $V_{\leq d-1}$, the subgraph $G[V_{\leq d-1}]$ is connected and contains C . Its vertices and edges lie in U (as they are not in H), and C borders the outer face of G , so $G[V_{\leq d-1}]$ and the outer face of G are connected within U , hence both lie in U_{out} .

Now let $v \in V_d$. Since $\text{depth}(v) = d \geq 1$, there exists $u \in V_{d-1}$ adjacent to v in G . The edge $\{v, u\}$ is not an edge of H , so it lies in U . Since $u \in V_{d-1} \subset U_{\text{out}}$ and $\{v, u\}$ is a connected subset of U containing u , the entire edge lies in U_{out} . The vertex v is an endpoint of this edge but is not in U , so v lies on the boundary of U_{out} , i.e., on the outer face of H . \square

Definition 2.7. Let G be a maximal planar graph with a plane embedding and outer cycle C . The *deep embedding* of G is the graph G' obtained from G by the following operation: for every neutral triangular face $\{u, v, w\}$ of G — including the outer face, whose vertices are the three vertices of C — add a new vertex x placed in that face and adjacent to each of u , v , and w . The vertex added inside the outer face is denoted x^* and called the *outer-cap vertex*; the three triangular faces it induces with the edges of C are the *outer-cap faces*. We henceforth view G' as embedded on the sphere S^2 , with no distinguished outer face.

Lemma 2.8. *Let G' be the deep embedding of a maximal planar graph G . Every face of G' is either an up triangle or a down triangle.*

Proof. We first establish that for any edge $\{p, q\}$ in G , the depths of p and q differ by at most 1. Suppose for contradiction that $\text{depth}(p) = d$ and $\text{depth}(q) = d + n$ for some $n \geq 2$. Since $\text{depth}(p) = d$, there exists a path of length d from p to some vertex of C . Prepending the edge $\{q, p\}$ gives a path of length $d + 1$ from q to C , so $\text{depth}(q) \leq d + 1 < d + n$, a contradiction. The case $\text{depth}(q) = d - n$ is handled identically: there exists a path of length $d - n$ from q to some vertex of C , and prepending the edge $\{p, q\}$ gives a path of length $d - n + 1 \leq d - 1 < d$ from p to C , contradicting $\text{depth}(p) = d$.

Since G is a triangulation, every interior face of G is a triangle $\{u, v, w\}$ with all three pairs adjacent. By the above, each pair of vertices in a triangle differs in depth by at most 1, so no triangle can contain vertices of depths d and $d + 2$ simultaneously. The possible depth patterns for a triangle in G are therefore exactly a neutral triangle, a down triangle, or an up triangle.

We now consider each case under the deep embedding.

Case 1: up triangle or down triangle. These triangles are not modified by the deep embedding, so they remain as faces of G' , satisfying the lemma.

Case 2: neutral triangle. The deep embedding inserts a new vertex x adjacent to u, v , and w , replacing the face $\{u, v, w\}$ with three new faces $\{u, v, x\}$, $\{v, w, x\}$, and $\{u, w, x\}$. It remains to determine the depth of x in G' . Since x is adjacent only to u, v , and w , every path in G' from x to C must pass through one of them, so x has strictly greater depth than u, v , and w . Each of the three new faces is thus a down triangle, satisfying the lemma. The same argument applies to the outer face: the outer-cap vertex x^* is adjacent to all three vertices of C (which lie at depth 0), so $\text{depth}(x^*) = 1$, and each of the three outer-cap faces is a down triangle.

Since every face of G' falls into one of these cases, the result follows. \square

3. QUADRILATERAL SEQUENCING

We now decompose the deep embedding into quadrilaterals by removing level edges, and define a deterministic sequence in which those quadrilaterals are visited.

Lemma 3.1. *Every interior face of G' has exactly one level edge.*

Proof. By the previous lemma, each interior face is an up triangle (depths $\{d, d + 1, d + 1\}$) or a down triangle (depths $\{d, d, d + 1\}$). In both cases, exactly one of the three vertex pairs has equal depth. \square

Lemma 3.2. *Let $e = \{p, q\}$ be any level edge of G' . Then e is the unique level edge of both faces incident to it.*

Proof. On the sphere, both faces T, T' incident to e are triangles. Since p and q have equal depth, e is a level edge of T and of T' , and by the previous lemma each has e as its unique level edge. \square

Definition 3.3. The *quadrilateral decomposition* of G' pairs each face of G' with the face on the other side of its (unique) level edge. Each pair, together with the four non-level edges of the two triangles, bounds a *quadrilateral* of the decomposition.

Remark 3.4. Because G' is taken on the sphere, every edge lies between two triangular faces, so the pairing above applies uniformly. In particular, each edge of C is

a level edge shared between one interior boundary down triangle (depths $\{0, 0, 1\}$, with the depth-1 vertex inside C') and one outer-cap down triangle (depths $\{0, 0, 1\}$, with apex x^*). The three resulting quadrilaterals, one per edge of C' , are the *boundary deep diamonds*; they are the outermost quadrilaterals of the decomposition.

Definition 3.5. Each quadrilateral is one of three types, classified by the depths of its two non-level vertices relative to the depth d of the shared level edge:

- a *shallow diamond*, formed by two up triangles, with vertex depths $(d - 1, d, d - 1, d)$ around the boundary;
- a *deep diamond*, formed by two down triangles, with vertex depths $(d + 1, d, d + 1, d)$ around the boundary;
- an *S quad*, formed by one up and one down triangle, with vertex depths $(d - 1, d, d + 1, d)$ around the boundary.

Remark 3.6. For the remainder of this section, fix a plane embedding of G' by designating one of the three outer-cap faces as the outer face of the plane drawing; the outer cycle of this plane embedding is the boundary of that designated outer-cap face. Orient this outer cycle counterclockwise so that the remaining faces of G' lie to its left. This induces a canonical cyclic order on the edges incident to each vertex and a notion of *left* and *right* on each triangle.

Definition 3.7. A *slice* of G' is a connected, simply connected region of the plane formed by the union of a subset of quadrilaterals from the decomposition, together with its closed boundary walk in G' .

Definition 3.8 (Move code). Each move is assigned a numerical code: anchor drop = 0, level add = 1, join = 2, ring completion = 3. Given a depth sequence Q_1, Q_2, \dots, Q_N , its *move-code string* is the word $m_2 m_3 \cdots m_N \in \{0, 1, 2, 3\}^{N-1}$, where m_n is the code of the move that produced Q_n from S_{n-1} .

Definition 3.9 (Initial quad). The depth sequence begins by choosing as Q_1 the boundary deep diamond whose resulting move-code string is lexicographically smallest among the three boundary deep diamonds. If multiple boundary deep diamonds yield the same move-code string, Q_1 is not uniquely determined; this case is called a *rotational tie* and corresponds to a symmetry of G' that permutes the boundary deep diamonds. The initial slice is $S_1 = Q_1$.

Remark 3.10. The tiebreak is recursive: the choice of Q_1 depends on the move-code strings produced by each of the three candidate starts, which in turn depend on the entire sequence each candidate produces. Equivalently, run the deterministic sequencing scheme from each of the three boundary deep diamonds and compare the resulting move-code strings; pick the start that produces the lexicographically smallest string.

Definition 3.11 (Anchor drop). Suppose a slice S_n has been constructed and the lower-rightmost portion of its boundary (in the fixed plane embedding) is a down triangle a whose right edge e is exposed. If there exists an S quad $Q \notin S_n$ whose up triangle b has e as its left edge, then the *anchor drop* sets

$$Q_{n+1} = Q, \quad S_{n+1} = S_n \cup Q.$$

The anchor drop introduces two new vertices: the second endpoint of b 's level edge (at depth one greater than a 's level edge) and the apex of Q 's down triangle (at depth two greater).

Definition 3.12 (Level add). Suppose a slice S_n has been constructed. Consider the quadrilaterals $Q \notin S_n$ such that exactly three of the four vertices of Q lie on the right boundary of S_n . Among these, the *level add* chooses the Q whose attachment to the right boundary occurs at the bottommost position (i.e., the last position encountered when scanning the right boundary from top to bottom). Then

$$Q_{n+1} = Q, \quad S_{n+1} = S_n \cup Q.$$

By construction, the level add introduces exactly one new vertex.

Definition 3.13 (Join). Suppose a slice S_n has been constructed. Consider the deep diamonds $Q \notin S_n$ such that one of the two down triangles comprising Q shares an edge with the right boundary of S_n (so two of Q 's four vertices lie on the right boundary). Among these, the *join* chooses the Q whose shared boundary edge is the bottommost in the right boundary scanned from top to bottom. Then

$$Q_{n+1} = Q, \quad S_{n+1} = S_n \cup Q.$$

Generically, the join introduces two new vertices: the second endpoint of Q 's level edge (depth d) and the apex of Q 's second down triangle (depth $d + 1$), where the shared boundary edge has depths $\{d, d + 1\}$.

Remark 3.14. Like the anchor drop, the join generically introduces two new vertices. Unlike the anchor drop, neither new vertex is at greater depth than the existing slice: the join extends the slice horizontally along the depth- d and depth- $(d + 1)$ rings rather than descending one level deeper.

Definition 3.15 (Ring completion). Suppose a slice S_n has been constructed. Consider the quadrilaterals $Q \notin S_n$ such that all four vertices of Q are already vertices of S_n . Among these, the *ring completion* chooses the Q whose attachment to the right boundary of S_n is bottommost (i.e., the last attachment encountered when scanning the right boundary from top to bottom). Then

$$Q_{n+1} = Q, \quad S_{n+1} = S_n \cup Q.$$

By construction, the ring completion introduces no new vertices.

Definition 3.16 (Move selection). At each step $n \geq 1$, the next quadrilateral Q_{n+1} is chosen by the first applicable move in the following order of precedence:

- (1) anchor drop;
- (2) level add;
- (3) join;
- (4) ring completion.

That is, each move is consulted only when no higher-precedence move applies.

Let N denote the total number of quadrilaterals in the decomposition of G' ; equivalently, $N = |F(G')|/2$, where $F(G')$ is the set of triangular faces of G' .

Theorem 3.17 (Termination and coverage). *The sequence Q_1, Q_2, \dots generated by repeatedly applying the move-selection rule starting from any choice of initial quadrilateral Q_1 terminates after exactly N steps. Moreover, every quadrilateral of the decomposition appears in the sequence exactly once, and $S_N = G'$.*

Proof. We prove the three claims comprising the theorem.

(1) *No quadrilateral is visited twice.* By construction, each move chooses $Q_{n+1} \notin S_n$ and sets $S_{n+1} = S_n \cup Q_{n+1}$. Hence $n \mapsto Q_n$ is injective.




FIGURE 1. The deep embedding G' of a small maximal planar graph (drawn with one outer-cap face as the outer face), with each quadrilateral Q_n of the canonical sequence labelled by its index and the move code (AD = anchor drop, LA = level add, J = join, RC = ring completion) of the move that produced it. Solid edges are non-level; dashed edges are level. Background colour encodes quadrilateral type: amber for shallow diamonds, teal for deep diamonds, pink for S quads. Outer-cycle vertices are blue, the outer-cap vertex x^* is red. The move-code string for this example is 01211333.

(2) *Each S_n is a slice.* We proceed by induction on n .

Base case. $S_1 = Q_1$ is a single quadrilateral, hence a closed topological disk on the sphere with boundary the closed walk along Q_1 's four perimeter edges.

Inductive step. Suppose S_n is a slice. We show that for each of the four moves, $S_{n+1} = S_n \cup Q_{n+1}$ is again a slice. Topologically, S_n is a closed disk and Q_{n+1} is a closed disk; their union is a closed disk iff their intersection $S_n \cap Q_{n+1}$ is a connected arc on each of their boundaries. We verify this in each case below by identifying the intersection precisely.

(2.1) *Anchor drop.* The new quadrilateral Q_{n+1} shares exactly one edge e with S_n : the right edge of the boundary down triangle a , which is also the left edge of the up triangle b . The intersection $S_n \cap Q_{n+1} = e$ is a single edge, a connected arc.

(2.2) *Level add.* Three vertices v_1, v_2, v_3 of Q_{n+1} lie on the right boundary of S_n . By the move's precedence, no anchor drop or higher-priority move applied at step $n + 1$. We claim the two perimeter edges of Q_{n+1} connecting these three vertices, namely $\{v_1, v_2\}$ and $\{v_2, v_3\}$, lie on the boundary of S_n . Suppose otherwise: then at least one of these edges has both adjacent faces in S_n^c , which means the face of G' on the side of that edge opposite Q_{n+1} is also outside S_n . By the structure of the move precedence and the inductive assumption, all such configurations would have been resolved by a higher-priority move first; hence at the moment level add fires, the only viable attachment configuration is the 2-edge path $\{v_1, v_2, v_3\}$. The intersection $S_n \cap Q_{n+1}$ is therefore this 2-edge path, a connected arc.

(2.3) *Join.* The deep diamond Q_{n+1} has one of its two down triangles sharing an edge e with the right boundary of S_n . The intersection $S_n \cap Q_{n+1} = e$, a single edge, is a connected arc. (The second down triangle of Q_{n+1} contributes two new vertices and lies entirely in S_n^c .)

(2.4) *Ring completion.* All four vertices of Q_{n+1} lie in $V(S_n)$, and by precedence none of the previous three moves applied. We claim that in this case, at least three of Q_{n+1} 's four perimeter edges lie on the boundary of S_n , and these edges together form a connected arc along the boundary of Q_{n+1} . Indeed, any perimeter edge of Q_{n+1} with both endpoints in S_n but neither side in S_n would force one of the higher-priority moves (specifically join or level add, applied with respect to the face on the other side of that edge); since no such move applied, the configuration is constrained so that the intersection $S_n \cap Q_{n+1}$ is a connected arc consisting of three or four of Q_{n+1} 's perimeter edges.

In all four cases the intersection $S_n \cap Q_{n+1}$ is a connected arc, so S_{n+1} is a closed topological disk and hence a slice. The boundary walk of S_{n+1} is obtained from that of S_n by deleting the arc $S_n \cap Q_{n+1}$ and inserting the complementary arc of Q_{n+1} 's boundary.

(3) *As long as $S_n \subsetneq G'$, some move applies.* Let S_n be a slice with $S_n \subsetneq G'$. The complement $G' \setminus S_n$ is a non-empty closed disk on the sphere whose boundary coincides with the boundary of S_n . Pick an edge e of the right boundary of S_n , and let F be the face of G' on the side of e opposite S_n ; then F lies in S_n^c . Let Q be the quadrilateral containing F .

Let k be the number of vertices of Q lying in $V(S_n)$, and let j be the number of perimeter edges of Q lying on the boundary of S_n . Since the edge e belongs to Q and lies on the boundary of S_n , we have $j \geq 1$ and $k \geq 2$.

Case $k = 2, j = 1$. Only the two endpoints of e are in S_n .

- If Q is an S quad and e is the left edge of Q 's up triangle, then writing a for the down triangle of S_n across e , the anchor drop hypothesis holds and the move applies (possibly with a different choice of a at the lower-rightmost position).
- If Q is a deep diamond, then e belongs to one of its down triangles, and the join hypothesis holds.
- If Q is a shallow diamond, then neither anchor drop nor join applies directly to Q . In this case, follow the boundary of S_n to find an adjacent face F' also in S_n^c whose containing quadrilateral Q' admits one of the four moves; such an adjacent quadrilateral exists because S_n^c is connected and is bounded by a closed walk in G' .

Case $k = 3$, $j = 2$. The three boundary vertices form a 2-edge path on Q 's outline, and the level add hypothesis holds. (No higher-priority move need have applied; if anchor drop also applies, the rule gives precedence to anchor drop.)

Case $k = 4$. All four vertices of Q are in $V(S_n)$, and $j \geq 1$ by assumption. If a higher-priority hypothesis (anchor drop, level add, join) holds, the corresponding move applies. Otherwise the ring completion hypothesis holds and that move applies.

In each case, some move adds either Q or an adjacent quadrilateral Q' to the slice, contradicting the assumption that no move applies. Hence as long as $S_n \subsetneq G'$, some move applies and S_{n+1} is well-defined.

Combining (1)–(3): the sequence strictly grows $|S_n|$ by exactly one quadrilateral per step, never revisits a quadrilateral, and must continue until $S_n = G'$. The total number of steps is therefore exactly N . \square

Remark 3.18. Two delicate sub-arguments in the proof of (2) and (3) deserve attention: (a) in (2.2) and (2.4), we relied on the move precedence to rule out configurations where the intersection $S_n \cap Q_{n+1}$ is disconnected; and (b) in (3), the shallow-diamond sub-case argues by “move to an adjacent face” but does not pin down which face. A more rigorous treatment would prove the equivalence of the following two statements: (i) the four moves cover every attachment configuration arising from a slice; (ii) at each step the move-selection rule produces a unique well-defined next quadrilateral.

REFERENCES

- [1] B. S. Baker, *Approximation algorithms for NP-complete problems on planar graphs*, Journal of the ACM, vol. 41, no. 1, pp. 153–180, 1994.