

PLANE DIAMOND COLORING

ERIC BAUERFELD

ABSTRACT.

NOTATION

For a coloring $C : V(G) \rightarrow S$ and a color $c \in S$, we write $C^{-1}(c) = \{v \in V(G) : C(v) = c\}$ for the preimage of c under C , i.e., the color class of c .

1. DEFINITIONS

Definition 1.1. Let G be a graph and let $u \in V(G)$. The *distance partition* of G from u is the partition $\{L_0, L_1, L_2, \dots\}$ of $V(G)$ obtained by breadth-first search from u :

$$L_0 = \{u\}, \quad L_{i+1} = \{v \in V(G) \setminus (L_0 \cup \dots \cup L_i) : v \text{ is adjacent to some } w \in L_i\}.$$

Equivalently, $L_i = \{v \in V(G) : d(v, u) = i\}$, where $d(v, u)$ denotes the graph distance between v and u in G . We call each L_i the *i-th level* of the partition.

Definition 1.2. Let G be a maximal planar graph. A *plane diamond coloring* of G is a proper 4-coloring C of G for which there exist a vertex $u \in V(G)$ and two distinct colors c_a, c_b such that, with respect to the distance partition $\{L_0, L_1, L_2, \dots\}$ of G from u ,

$$C^{-1}(c_a) \subseteq \bigcup_{i \text{ even}} L_i \quad \text{and} \quad C^{-1}(c_b) \subseteq \bigcup_{i \text{ odd}} L_i.$$

2. RESULTS

Remark 2.1. Definition 1.2 imposes a structural condition on 4-colorings of maximal planar graphs strictly stronger than the conclusion of the Four Color Theorem [1, 2]: it requires not merely the existence of a proper 4-coloring, but the existence of a proper 4-coloring together with a root u such that two of the four color classes are separated by the parity of the BFS layering from u .

Conjecture 2.2. *Every maximal planar graph G has a plane diamond coloring.*

Theorem 2.3. *The preceding conjecture is false. Moreover, the smallest counterexample has order 13, and is unique up to isomorphism among triangulations of order at most 13.*

Proof. Let G be the maximal planar graph on 13 vertices with graph6 string¹

LQh0?SFIwnTYTi

2010 *Mathematics Subject Classification.* Primary .

¹We use the standard graph6 encoding of McKay; see [3].

shown in Figure 1. Equivalently, G has edge set

$$\begin{aligned} &\{0, 2\}, \{0, 4\}, \{0, 11\}, \{1, 3\}, \{1, 5\}, \{1, 12\}, \{2, 4\}, \{2, 9\}, \{2, 11\}, \\ &\{3, 5\}, \{3, 10\}, \{3, 12\}, \{4, 7\}, \{4, 9\}, \{4, 11\}, \{5, 8\}, \{5, 10\}, \{5, 12\}, \\ &\{6, 7\}, \{6, 8\}, \{6, 9\}, \{6, 10\}, \{6, 11\}, \{6, 12\}, \{7, 8\}, \{7, 9\}, \{7, 10\}, \\ &\{7, 11\}, \{8, 9\}, \{8, 10\}, \{8, 12\}, \{9, 11\}, \{10, 12\}. \end{aligned}$$

We have $|V(G)| = 13$ and $|E(G)| = 33 = 3 \cdot 13 - 6$, so G is a triangulation.

By Definition 1.2, it suffices to show that for every root $u \in V(G)$, no proper 4-coloring C of G admits two distinct colors c_a, c_b with $C^{-1}(c_a)$ contained in the union of even-indexed levels and $C^{-1}(c_b)$ contained in the union of odd-indexed levels of the distance partition from u .

For a fixed root u , the existence of such a triple (C, c_a, c_b) is equivalent to 4-colorability of the auxiliary graph H_u obtained from G by adjoining two new vertices α, β , joining α to every vertex in odd-indexed levels, joining β to every vertex in even-indexed levels, and adding the edge $\{\alpha, \beta\}$. Indeed, in any proper 4-coloring of H_u the colors of α and β are distinct and absent from the odd-parity and even-parity layers of G respectively, yielding $c_a := C(\alpha)$ and $c_b := C(\beta)$. Conversely, given a 4-coloring satisfying the parity-separation condition, setting $C(\alpha) := c_a$ and $C(\beta) := c_b$ extends it to a proper 4-coloring of H_u .

A direct computation (using Sage's `chromatic_number`) verifies that $\chi(H_u) > 4$ for every $u \in V(G)$, so G admits no plane diamond coloring.

For minimality and uniqueness, we exhaustively enumerated every maximal planar graph of order at most 13 using Sage's `graphs.planar_graphs` generator (with `minimum_connectivity=3` and `maximum_face_size=3`). The numbers of triangulations 1, 1, 2, 5, 14, 50, 233, 1249, 7595, 49566 at orders 4, 5, \dots , 13 respectively (matching OEIS A000109) were each tested for the existence of a plane diamond coloring, and exactly one — the graph G above, occurring at order 13 — was found to lack one. \square




FIGURE 1. The unique smallest maximal planar graph with no plane diamond coloring; it has 13 vertices and degree sequence $(6, 6, 6, 6, 6, 6, 6, 5, 5, 4, 4, 3, 3)$.

REFERENCES

- [1] K. Appel and W. Haken, *Every planar map is four colorable*, Illinois Journal of Mathematics, vol. 21, no. 3, pp. 429–567, 1977.
- [2] N. Robertson, D. Sanders, P. Seymour, and R. Thomas, *The four-colour theorem*, Journal of Combinatorial Theory, Series B, vol. 70, no. 1, pp. 2–44, 1997.
- [3] B. D. McKay, *Description of graph6, sparse6 and digraph6 encodings*, <https://users.cecs.anu.edu.au/~bdm/data/formats.txt>.