

COLORED EDGE FLIP CLASSES

ERIC BAUERFELD

ABSTRACT.

1. MOTIVATION

The Four Color Theorem asserts that every planar graph is properly 4-colorable, or equivalently that no maximal planar graph G satisfies $\chi(G) \geq 5$. Suppose, towards a contradiction, that such a graph exists; let G_0 be one of minimum order. Any structural property shared by every maximal planar graph H with $|V(H)| = |V(G_0)|$ is then automatically inherited by G_0 , and any property *not* satisfied by G_0 excludes a portion of the class of maximal planar graphs from playing the role of a minimum counterexample.

This paper investigates one such property: behavior under an edge flip. Our principal observation (Theorem 4.4) is that every edge flip of a minimum-order 5-chromatic maximal planar graph yields a 4-colorable graph. In particular, no such graph is *flip-symmetric*, where we call a maximal planar graph G flip-symmetric when some admissible flip at an edge of G returns a graph isomorphic to G . The search for a counterexample to the Four Color Theorem may therefore be confined to the complement of the class \mathcal{F} of flip-symmetric graphs.

2. PRELIMINARIES

Let G be a maximal planar graph with $|V(G)| \geq 4$, embedded in the plane so that every face — including the outer face — is a triangle. Every edge $uv \in E(G)$ is then shared by exactly two triangular faces uvw and uvx whose union is a quadrilateral $uwvx$ with diagonal uv .

Definition 2.1 (Edge flip). Let G be a maximal planar graph and let $uv \in E(G)$ be an edge whose two incident triangular faces are uvw and uvx . The *edge flip* (or *diagonal flip*) at uv is the operation that deletes the edge uv and inserts the edge wx in its place, replacing the two triangles uvw and uvx by the two triangles uwv and vwx . The flip is *admissible* if $wx \notin E(G)$; otherwise the resulting multigraph is not simple and the flip is forbidden.

3. FLIP-SYMMETRIC MAXIMAL PLANAR GRAPHS

For a maximal planar graph G and an admissible edge $uv \in E(G)$ with incident triangles uvw , uvx , write

$$G^{\text{flip}(uv)} = (V(G), (E(G) \setminus \{uv\}) \cup \{wx\})$$

for the graph obtained from G by flipping uv .




FIGURE 1. An edge flip replaces the diagonal uv of the quadrilateral $uwvx$ with the diagonal wx .

Definition 3.1 (Flip-symmetric graph). A maximal planar graph G is *flip-symmetric* if there exists an admissible edge $uv \in E(G)$ such that $G^{\text{flip}(uv)} \cong G$. We write \mathcal{F} for the class of flip-symmetric maximal planar graphs.

Definition 3.2 (Colored edge flip class). Let G be a maximal planar graph and let φ be a proper 4-coloring of G . The *colored edge flip class* of (G, φ) is the set

$$\mathcal{C}(G, \varphi) = \{ G^{\text{flip}(uv)} : uv \in E(G), \text{ the flip at } uv \text{ is admissible, and } \varphi(w) \neq \varphi(x) \},$$

where w, x are the third vertices of the two triangular faces of G containing uv . Equivalently, $\mathcal{C}(G, \varphi)$ is the set of graphs obtained from G by an admissible edge flip under which φ remains a proper 4-coloring.

4. A MINIMAL FOUR-COLORABLE COUNTEREXAMPLE

Definition 4.1 (Edge-deletion subgraph). Let G be a maximal planar graph and $uv \in E(G)$. The *edge-deletion subgraph at uv* is the spanning subgraph $G - uv = (V(G), E(G) \setminus \{uv\})$. Write $\mathcal{D}(G) = \{G - uv : uv \in E(G)\}$.

Lemma 4.2. *Let G_0 be a maximal planar graph of minimum order with $\chi(G_0) \geq 5$. Then every $H \in \mathcal{D}(G_0)$ is 4-colorable.*

Proof. Fix $uv \in E(G_0)$ and let G_0/uv denote the simple planar graph obtained by contracting uv and discarding parallel edges. Since $|V(G_0/uv)| = |V(G_0)| - 1$, the minimality of G_0 supplies a proper 4-coloring c of G_0/uv . Let z be the contracted vertex and define $c' : V(G_0) \rightarrow \{1, 2, 3, 4\}$ by $c'(u) = c'(v) = c(z)$ and $c'(y) = c(y)$ for $y \notin \{u, v\}$. Every edge of $G_0 - uv$ is either disjoint from $\{u, v\}$ or incident to exactly one of them; in either case the corresponding edge of G_0/uv has distinct endpoints under c , so c' assigns its endpoints distinct colors. The edge uv itself is absent from $G_0 - uv$, so c' is a proper 4-coloring of $G_0 - uv$. \square

Lemma 4.3. *Let G_0 be a maximal planar graph of minimum order with $\chi(G_0) \geq 5$, fix $uv \in E(G_0)$, and let φ be any proper 4-coloring of $G_0 - uv$. Write $a = \varphi(u)$ and let b, c, d denote the three remaining colors. Then:*

- (1) $\varphi(v) = a$;
- (2) *the subgraph of $G_0 - uv$ induced by the vertices of color a or b contains a path from u to v ;*

- (3) *the subgraph of $G_0 - uv$ induced by the vertices of color a or c contains a path from u to v .*

Proof. (1) If $\varphi(v) \neq a$ then φ is already a proper 4-coloring of G_0 , since the only edge of G_0 absent from $G_0 - uv$ is uv and its endpoints have distinct colors. This contradicts $\chi(G_0) \geq 5$, so $\varphi(v) = a$.

(2) Suppose, for contradiction, that u and v lie in distinct connected components of the subgraph of $G_0 - uv$ induced by the color classes a and b . Let C be the component containing u , and define $\varphi': V(G_0) \rightarrow \{a, b, c, d\}$ by swapping colors $a \leftrightarrow b$ on C and leaving every other vertex unchanged. Then φ' is a proper 4-coloring of $G_0 - uv$ with $\varphi'(u) = b$ and $\varphi'(v) = a$, contradicting part (1) applied to φ' .

(3) Identical to (2) with c in place of b . \square

Theorem 4.4. *Let G be a minimum-order maximal planar graph with $\chi(G) \geq 5$. Then for every edge $e \in E(G)$, the graph induced by an edge flip of e is 4-colorable.*

Proof. Fix an edge $e = uv \in E(G)$, and let F_0, F_1 be the two triangular faces of G incident to e , so that $\{w, x\} = (V(F_0) \cup V(F_1)) \setminus \{u, v\}$. By Lemma 4.2, $G - e$ admits a proper 4-coloring φ .

Case 1: $\varphi(w) \neq \varphi(x)$. Then φ is also a proper 4-coloring of the graph induced by the edge flip of e .

Case 2: $\varphi(w) = \varphi(x)$. Set $a = \varphi(u)$; by Lemma 4.3(1), $\varphi(v) = a$ as well, and the edges $uw, vw \in E(G - e)$ force $\varphi(w) \neq a$. Choose a color $b \notin \{a, \varphi(w)\}$. By Lemma 4.3, there is a path P from u to v in the subgraph of $G - e$ induced by the vertices of color a or b . Let $\{c, d\} = \{1, 2, 3, 4\} \setminus \{a, b\}$; then $\varphi(w) = \varphi(x) \in \{c, d\}$.

Any path from w to x in the subgraph of $G - e$ induced by the vertices of color c or d would, in the plane embedding of $G - e$, cross P ; but its vertices have colors in $\{c, d\}$ and the vertices of P have colors in $\{a, b\}$, and these sets are disjoint, so the two paths share no vertex. Hence w and x lie in distinct connected components of the $\{c, d\}$ -colored subgraph of $G - e$. Swapping colors $c \leftrightarrow d$ on the component containing w yields a proper 4-coloring of $G - e$ in which $\varphi(w) \neq \varphi(x)$, reducing to Case 1. \square




FIGURE 2. Case 2 of the proof of Theorem 4.4: u, v share color a and w, x share color c . The $\{a, b\}$ -Kempe path P from u to v separates w from x in the plane, so no $\{c, d\}$ -path between w and x can avoid crossing P ; since the color sets $\{a, b\}$ and $\{c, d\}$ are disjoint, no such path exists.